

Animal Health Foundation Blog

Dr. Jean Dodds – Dog Vaccine Protocol

The following vaccine protocol is offered for those dogs where minimal vaccinations are advisable or desirable. The schedule is one I recommend and should not be interpreted to mean that other protocols recommended by a veterinarian would be less satisfactory. It's a matter of professional judgment and choice.

9 – 10 weeks of age

Distemper + Parvovirus, MLV
e.g. Merck Nobivac (Intervet Progard) Puppy DPV

14 – 15 weeks of age

Distemper + Parvovirus, MLV

18 weeks of age

Parvovirus only, MLV

Note: New research states that last puppy parvovirus vaccine should be at 18 weeks old.

20 weeks or older, if allowable by law

Rabies – give 3-4 weeks apart from other vaccines
Mercury-free (thimerosal-free, TF)

1 year old

Distemper + Parvovirus, MLV

This is an optional booster or titer. If the client intends *not* to booster after this optional booster or intends *to* retest titers in another three years, this optional booster at puberty is wise.

1 year old

Rabies – give 3-4 weeks apart from other vaccines
3-year product if allowable by law; mercury-free (TF)

Perform vaccine antibody titers for distemper and parvovirus every three years thereafter, or more often, if desired. Vaccinate for rabies virus according to the law,

except where circumstances indicate that a written waiver needs to be obtained from the primary care veterinarian. In that case, a rabies antibody titer can also be performed to accompany the waiver request. Visit [The Rabies Challenge Fund](#) for more information.

W. Jean Dodds, DVM
[Hemopet](#) / [NutriScan](#)
11561 Salinaz Avenue
Garden Grove, CA 92843

This entry was posted on Saturday, December 9th, 2017 at 12:46 am and is filed under [Dogs](#), [Healthy Pets](#).
